arXiv:hep-ph/0409200v1 16 Sep 2004

PSI-PR-04-10
hep-ph/0409200

# SDECAY - A Fortran Code For SUSY Particle Decays In The MSSM

M. M. MÜHLLEITNER
*Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland*

The Fortran code `SDECAY` calculates the decay widths and branching ratios of all supersymmetric (SUSY) particles in the Minimal Supersymmetric Standard Model (MSSM), including higher order effects. The usual two-body decays of sfermions and gauginos as well as the three-body decay modes of charginos, neutralinos and gluinos are included. Furthermore, the three-body stop and sbottom decays and even the four-body stop decays are calculated. The important loop-induced decays, the QCD corrections to the two-body widths involving strongly interacting particles and the dominant electroweak effects to all processes are evaluated as well.

## 1 Introduction

The search for SUSY particles is a major goal at present and future colliders. Once they are found, their properties are expected to be determined with an accuracy of a few per cent at the LHC and at the per cent level and below at future $e^+e^-$ linear colliders. To match the expected experimental accuracy, programs are needed for the calculation of the SUSY particle spectrum, of the production cross sections and of the SUSY particle decay widths and branching ratios with high precision, also including higher order effects. The Fortran code `SDECAY`[1] which is presented in the following deals with the decay of SUSY particles in the MSSM and includes the most important higher order effects. Up to the user's choice, the mass spectrum and soft SUSY breaking parameters are obtained from the renormalization group evolution program `SuSpect`[2] or from an input file in the SUSY Les Houches Accord (SLHA) format[3]. `SDECAY` then evaluates the various couplings of the SUSY particles and MSSM Higgs bosons and calculates the decay widths and branching ratios.[a]

## 2 The tree-level and QCD corrected two-body decays

The following tree-level two-body sfermion decays have been implemented in the program[b]

$$\tilde{f}_i \;\to\; \tilde{\chi}_j f \qquad \tilde{\chi}_j : \text{ chargino, neutralino}$$

---

[a] For details of the implementation of the MSSM see the user's manual[1].

[b] Here and in the following the distinction between the two isospin (s)fermion partners is discarded.

$$\begin{aligned} \tilde{f}_i &\to \tilde{f}_j V \qquad V:\ W, Z \\ \tilde{f}_i &\to \tilde{f}_j \Phi \qquad \Phi:\ h, H, A, H^{\pm} \end{aligned} \tag{1}$$

The included two-body decays for the squarks, the gluino, the charginos and neutralinos are

$$\tilde{q}_i \to \tilde{g}\, q, \qquad \tilde{g} \to \tilde{q}_i q \tag{2}$$

$$\tilde{\chi}_i \to \tilde{\chi}_j V, \qquad \tilde{\chi}_i \to \tilde{\chi}_j \Phi \quad \text{and} \quad \tilde{\chi}_i \to \tilde{f}_j f \tag{3}$$

In case of the gauge mediated SUSY breaking model, the following 2-body decays into the lightest SUSY particle, the gravitino, have been implemented[1]

$$\tilde{\chi}_1^0 \to \tilde{G}\gamma, \tilde{G}Z, \tilde{G}\phi \ \ (\phi = h, H, A) \quad \text{and} \quad \tilde{\tau}_1 \to \tilde{G}\tau \tag{4}$$

A new feature compared to many existing decay programs, is the inclusion of the SUSY-QCD corrections to 2-body decays involving coloured particles[1]:

$$\tilde{q}_i \to \tilde{\chi}_j q, \quad \tilde{q}_i \to \tilde{q}_j V, \quad \tilde{q}_i \to \tilde{q}_j \Phi, \quad \tilde{q}_i \to \tilde{g} q \tag{5}$$

$$\tilde{g} \to \tilde{q}_i q, \quad \tilde{\chi}_i \to \tilde{q}_j q \tag{6}$$

We are taking running parameters at the scale of the electroweak symmetry breaking for the gauge and third generation Yukawa couplings as well as the soft SUSY-breaking parameters and third-generation sfermion mixing angles. In this way, the bulk of the electroweak corrections has been taken into account. Finally, `SDECAY` also evaluates the top quark decays in the MSSM:

$$t \to bW^+, \quad t \to bH^+ \quad \text{and} \quad t \to \tilde{t}_1 \tilde{\chi}_1^0 \tag{7}$$

## 3 The loop-induced and the multibody decays

If the 2-body decays are closed, multibody decays will be dominant. The implemented gaugino, gluino, stop and sbottom three-body decays[1] are

$$\tilde{\chi}_i \to \tilde{\chi}_j f \bar{f}, \qquad \tilde{\chi}_i \to \tilde{g} q \bar{q} \tag{8}$$

$$\tilde{g} \to \tilde{\chi}_i q \bar{q}, \qquad \tilde{g} \to \tilde{t}_1 \bar{b} W^-, \quad \tilde{g} \to \tilde{t}_1 \bar{b} H^- \tag{9}$$

$$\tilde{t}_i \to b W^+ \tilde{\chi}_1^0, \quad \tilde{t}_i \to b H^+ \tilde{\chi}_1^0 \tag{10}$$

$$\tilde{t}_i \to b l^+ \tilde{\nu}_l, \quad \tilde{t}_i \to b \tilde{l}^+ \nu_l, \quad \tilde{t}_i \to \tilde{b}_j f \bar{f}, \quad \tilde{t}_2 \to \tilde{t}_1 f \bar{f} \tag{11}$$

$$\tilde{b}_i \to t l^- \tilde{\nu}_l^*, \quad \tilde{b}_i \to t \tilde{l}^- \bar{\nu}_l, \quad \tilde{b}_i \to \tilde{t}_j f \bar{f}, \quad \tilde{b}_2 \to \tilde{b}_1 f \bar{f} \tag{12}$$

In the 3-body decays, the mixing of the third-generation sfermions as well as the final-state fermion masses have been taken into account[1,4]. Moreover, loop-induced decays of the lightest stop, gluino and the next-to-lightest neutralino[1] might become important and have therefore been included:

$$\tilde{\chi}_2^0 \to \tilde{\chi}_1^0 \gamma, \qquad \tilde{g} \to \tilde{\chi}_1^0 g \qquad \text{and} \qquad \tilde{t}_1 \to \tilde{\chi}_1^0 c \tag{13}$$

If the 3-body decays are kinematically forbidden, the following $\tilde{t}_1$ 4-body decay can compete with the loop-induced $\tilde{t}_1$ decay and has been implemented:

$$\tilde{t}_1 \quad \rightarrow \quad b\tilde{\chi}_1^0 f\bar{f} \tag{14}$$

## 4 Structure of `SDECAY`

Besides the necessary `SuSpect` files, the program `SDECAY` is composed of 3 files: (i) `sdecay.in`, where the user can choose various options, such as whether or not to include the QCD corrections to 2-body decays, the multibody and/or loop decays, the GMSB decays, the top decays. One can choose the scale and the number of loops for the running couplings and also, where to take the spectrum from, *i.e.* from `SuSpect` or any input file in the SLHA format; (ii) `sdecay.f` is the main body of the code. It evaluates the various couplings and calculates the decay widths and branching ratios; (iii) `sdecay.out`, which gives out the branching ratios and total widths, either in a simple transparent form or in the SLHA format, which is up to the user's choice. In addition, the whole SUSY spectrum, mixing matrices and further parameters used for the calculation of the couplings and decay widths are written out. The necessary files and the makefile for the compilation can be found at the `SDECAY` homepage:
http://people.web.psi.ch/muehlleitner/SDECAY/.

## 5 Summary

The Fortran code `SDECAY` calculates all SUSY particle 2-body tree-level and QCD corrected decays, 3-body and loop-decays as well as $\tilde{t}_1$ 4-body and top-decays. The program is rather fast, flexible and maintained regularly to include upgrades and newest theoretical developments as well as experimental needs.

## References

1. M. Mühlleitner, A. Djouadi and Y. Mambrini, arXiv:hep-ph/0311167 [and references therein].
2. A. Djouadi, J.-L. Kneur and G. Moultaka, arXiv:hep-ph/0211331.
3. P. Skands et al., JHEP **0407** (2004) 036 [arXiv:hep-ph/0311123].
4. A. Djouadi, Y. Mambrini and M. Mühlleitner, Eur. Phys. J. **C20** (2001) 563 [arXiv:hep-ph/0104115].